



Western SARE

Phil Rasmussen, Coordinator
Utah State University
Agricultural Science Building
Room 305
4865 Old Main Hill
Logan, Utah 84322-4865
phone: (435) 797-2257
fax: (435) 797-3344

Professional Development Program

Morgan Doran
California PDP Coordinator
Livestock & Natural Resource Farm
Advisor
University of California Cooperative
Extension
501 Texas Street
Fairfield, CA 94533
707.784.1326
mpdor@ucdavis.edu

Western SARE Grant Categories

- Research & Education
- Professional Development
- Farmer/Rancher
- Professional + Producer
- Graduate Student
- Sustainable Farm Tours

Go to <http://wsare.usu.edu>
Click on: Apply for a Grant



TRIPLE-CROPPING DAIRY FORAGE

Situation

Excess nitrate in ground-water under California dairies is common, and recent regulations prohibit applications

Research and Education Grant

Title: Triple-Cropping Dairy Forage Production Systems through Conservation Tillage in California's San Joaquin Valley

Project Number: SW08-060

Principal Investigator

Jeff Mitchell
Department of Plant Sciences
University of California
Kearny Agricultural Center
9240 St. Riverbend Avenue
Parlier, CA 93648
559.303.9689
Mitchell@uckac.edu

Participants

Marsha Campbell-Mathews
UC Cooperative Extension
Stanislaus County
David Crohn
UC Department of Environmental Sciences
Riverside
Carol Frate
UC Cooperative Extension
Tulare County
William Horwath
UC Department of Land, Air, and Water Resources
Davis
Anil Shrestha
Department of Plant Science and Mechanized Agriculture
California State University
Fresno
Danny Petersen
Aldon Petersen and Sons
Turlock
Frank Gwerder
Gwerder Dairy
Dino Giacomazzi
Giacomazzi Dairy
SARE Grant: \$118,100



Dairy farmer and project partner Frank Gwerder operates the tractor pulling the no-till drill planting triple-crop sudangrass following strip-till silage corn at Gwerder Dairy in Modesto. – Photos by Jeff Mitchell

that exceed 140% of crop removal. This standard could force many dairies out of business, especially small ones, if they cannot acquire additional land.

Triple-cropping – growing three forage crops a year –



No-till established sudangrass triple-crop emerges in corn residue at Gwerder Dairy.

utilizes more manure nutrients on the same ground and provides more feed. But it depends on timely harvests of winter small grain forages and summer corn silage.

Conservation tillage practices, used in other parts of the United States but essentially new in California, eliminate intercrop tillage operations, shorten the cropping interval, and allow for faster planting and harvest. This could make triple-cropping a potentially reliable component of sustainable nutrient management plans.

Objectives

1. Increase the reliability of triple-cropping dairy forage production with the use of conservation tillage practices as a means of increasing forage biomass and nutrient uptake by deter-

SARE's mission is to advance—to the whole of American agriculture—innovations that improve profitability, stewardship, and quality of life by investing in groundbreaking research and education.

The Western Region, one of four SARE regions nationwide, is administered through Utah State University.

Western SARE:
<http://wsare.usu.edu>

National SARE
www.sare.org

TRIPLE-CROPPING DAIRY FORAGE

mining production rates and N removal in triple-cropped forage fields compared with standard double-cropped fields

2. Evaluate triple-cropping compared with standard double-cropping forage production in terms of whole dairy nitrogen budgets and profitability
3. Extend widely information developed by the project to dairy farmers, consultants, and industry groups via a variety of extension education means
4. Track changes in the adoption of conservation tillage forage production practices in the San Joaquin Valley as evidenced by conservation tillage acreage surveys conducted by California's Conservation Tillage Workgroup and conservation tillage equipment sales records

Actions

Field evaluations of triple cropping (the production of three forage crops in a given year versus standard double-cropping practices) have been established at the San Joaquin Valley dairies of Frank Gwerder in Modesto and Danny Peterson in Turlock.

In the first year of the project, these activities have been conducted:

- Biomass production and tissue nitrogen content for each of the crops in these dairy silage fields have been sampled and determined at approximately weekly intervals
- Soil nitrogen has been monitored at key times during each crop cycle



At right, strip-till planted silage corn compares with conventionally planted silage corn using full intercrop tillage at left.

- Air temperature information has been collected
- Large-scale field yield determinations have been made in conjunction with each partner farmer
- Data from the monitoring activities are being compiled and summarized

An adjunct evaluation of twin-row strip-tilled corn vs. single-row strip-tilled corn was initiated at the dairy of Dino Giacomazzi in Hanford.

The project team also hosted public seminars featuring John Landers, a noted no-till expert in Brazil, at the dairy farms of Frank Gwerder in Modesto and Tom Barcellos in Tipton.

Information related to the goals of the SARE project and initial experiences were presented at the Conservation Tillage Workgroup's display during the 2009 World Ag Expo at Tulare in February.

Results

During the study's first year, the project team noted two potentially significant constraints of conservation

tillage practices used to facilitate triple-cropping at the dairies:

At the Petersen dairy, manganese deficiency occurred in the strip-till triple-cropped corn crop and reduced silage yield. It was hypothesized that this resulted from nutrient stratification in the soil of the conservation tillage system. To address the situation, a shallow disking was done before corn planting in 2009.

At the Gwerder dairy, problems aligning the corn planter following strip-tilling were observed, resulting in reduced forage production in the conservation tillage system. The situation has been discussed with Frank Gwerder, and plans were made to remedy the problem in the next corn planting.

Work Remaining

In the coming cycle, the project team will continue field monitoring and consolidating the data and applying it to the APTRAC Manure N Management Model.

The project team will also target more extension education events for San Joaquin Valley dairy producers.