

Characteristics	Site A	Site B	Site C	Site D
Location	Waterside	Waterside	Thalassia	Open Bay
Organic	Waterside	Organic	Organic	Open Bay
Crop in adjacent field	Wheat	Coke crops, seaweed	Corn, pepper, cotton, tomatoes, leeks	Coke crops, cotton
Year bigholgers were collected	2000	2000	2001	1999
Wind direction	S-E-W	S-W	W-E	W-E
Bigholger length (in 1972)	W-910	W-910	W-910	W-910
Plant studied	<i>A. subulifolius</i> <i>R. pulchellus</i> <i>C. graminea</i> <i>C. Ray-Hammett</i> <i>R. arifolius</i> <i>R. confertifolia</i>	<i>A. subulifolius</i> <i>A. multiflorus</i> <i>A. multiflorus</i> <i>C. Ray-Hammett</i> <i>A. graminea</i> <i>R. confertifolia</i>	<i>A. subulifolius</i> <i>A. multiflorus</i> <i>A. multiflorus</i> <i>C. Ray-Hammett</i> <i>C. graminea</i> <i>R. confertifolia</i>	<i>A. subulifolius</i> <i>A. multiflorus</i> <i>A. multiflorus</i> <i>C. graminea</i> <i>A. multiflorus</i> <i>R. confertifolia</i>
Total number of bigholger species	13	35	13	13
% Crops	<10%	30-51%	30-10%	50-10%