

Grazing Riparian Ecosystems: Season of Use

Miranda A. Meehan
Research Assistant

Kevin K. Sedivec
Extension Rangeland Management
Specialist

Edward S. DeKeyser
Associate Professor

Livestock often are attracted to riparian systems for the availability of forage and water, shade and smooth terrain.

Riparian ecosystems are extremely productive, with some providing 81 percent of the summer forage utilized by grazing livestock (Roath and Krueger 1982). Changes in riparian vegetation induced by grazing can result in a decline in soil health, loss of biotic diversity, degradation of wildlife habitat, reduced water quality and alterations in stream hydrology.

However, grazing has been found to be important for the proper functioning of many riparian zones. Implementation of proper grazing management practices is essential to prevent degradation by livestock and improve riparian health. The season of use or timing is critical when designing a grazing system aimed at maintaining healthy riparian systems.

Spring

Spring grazing is an effective management strategy for utilizing riparian zones. Upland forages are more palatable in the spring, creating a grazing behavior pattern in which livestock select upland sites, using riparian areas less frequently at this time. This results in a more uniform

grazing distribution pattern (Parsons et al. 2003, DelCurto et al. 2005).

In the spring, upland plant species have high forage quality and are low in dry matter. Crude protein and total digestible nutrients are at their highest and fiber content lowest in the spring. Crude protein and digestibility reduce into the summer period as the plant matures and fiber increases. Riparian ecosystems grazed in the spring have greater potential for regrowth, compared with summer use, resulting in greater production and protection of stream banks.

One disadvantage of spring use of riparian ecosystems is an increased chance of soil compaction and erosion due to increased soil moisture during this period.



Summer


Summer grazing is second only to seasonlong grazing as having the most negative impact on the health of riparian vegetation. Summer grazing of riparian ecosystems results in greater utilization of riparian areas than adjacent uplands because livestock are attracted to water, cooler temperatures, shade for fly control and higher-quality vegetation. Summer vegetation removal has less potential for regrowth, decreasing production and cover potential, leaving stream banks more exposed to peak flows, and increasing erosion and sediment loads in the stream.



Fall/Dormant Season

Although most research favors spring utilization of riparian pastures for maintenance of riparian health, fall grazing can be beneficial to riparian health. Riparian vegetation is more palatable and of greater nutritive quality than adjacent upland vegetation late in the growing season, with many sedge species outranking key upland forage species in crude protein and energy content (Kauffman et al. 1983).

Dormant-season grazing reduces the impacts of trampling on soil compaction because river levels are generally low and flood plains are either dry or frozen (Sedgwick and Knopf 1991). Overuse of riparian areas in the dormant season leads to increased utilization of shrubs and trees (Clary and Leininger 2000). However, these impacts can be reduced by careful monitoring and controlling the time livestock are present and grazing distribution by using other management practices.



Conclusions

Implementation of proper grazing management practices is required to prevent overgrazing by livestock and improve riparian health. The season of use or timing is critical when designing a grazing system aimed at maintaining the health of riparian systems (Table 1). Spring and dormant-season grazing have been shown to be effective for livestock utilizing riparian zones.

Table 1. Impact of season of grazing on upland and riparian vegetation communities in the northern Plains.

Season	Spring	Summer	Fall
Upland Vegetation	-	+	-
Riparian Vegetation	+	-	0
	- Negative	+ Positive	0 Neutral

Literature Cited

- Clary, W.P., and W.C. Leininger. 2000. Stubble height as a tool for management of riparian areas. *J. Range Manage.* 53(6):562-573.
- DelCurto, T., M. Porath, C.T. Parsons and J.A. Morrison. 2005. Management strategies for sustainable beef cattle grazing on forested rangelands in the Pacific Northwest. *Rangeland Ecol. & Manage.* 58(2): 119-127.
- Kauffman, J.B., W.C. Krueger and M. Vavra. 1983. Effects of late season cattle grazing on riparian plant communities. *J. Range Manage.* 36(6): 685-691.
- Parsons, C.T., P.A. Momont, T. Delcurto, M. McInnis and M.L. Porath. 2003. Cattle distribution patterns and vegetation use in mountain riparian areas. *J. Range Manage.* 56(4): 334-341.
- Roath, L.R., and W.C. Kruegar. 1982. Cattle grazing influence on a mountain riparian zone. *J. Range Manage.* 35(1): 100-103.
- Sedgwick, J.A., and F.L. Knopf. 1991. Prescribed grazing as a secondary impact in a western riparian floodplain. *J. Range Manage.* 44(4): 369-373.

Other publications in this series

- R-1539 [Riparian Ecosystems of North Dakota](#)
- R-1540 [Grazing Riparian Ecosystems: Grazing Systems](#)
- R-1541 [Grazing Riparian Ecosystems: Grazing Intensity](#)
- R-1543 [Grazing Riparian Ecosystems: Water Developments](#)

For more information on this and other topics, see: www.ag.ndsu.edu

NDSU encourages you to use and share this content, but please do so under the conditions of our Creative Commons license. You may copy, distribute, transmit and adapt this work as long as you give full attribution, don't use the work for commercial purposes and share your resulting work similarly. For more information, visit www.ag.ndsu.edu/agcomm/creative-commons.

North Dakota State University does not discriminate on the basis of age, color, disability, gender identity, marital status, national origin, public assistance status, sex, sexual orientation, status as a U.S. veteran, race or religion. Direct inquiries to the Vice President for Equity, Diversity and Global Outreach, 205 Old Main, (701) 231-7708.

County Commissions, NDSU and U.S. Department of Agriculture Cooperating.

This publication will be made available in alternative formats for people with disabilities upon request, (701) 231-7881.